## JOURNAL OF CARDIOVASCULAR, NEUROVASCULAR & STROKE

	Contents	
Volum	ne 6 Number 1 N	March 2024
No.	Title	Page
Case R	leport	
1.	CHALLENGING CATHETER MANAGEMENT IN A RARE C	CASE OF
	MYELOMENINGOCELE WITH SITUS INVERSUS TOTALIS:	A CASE
	REPORT	
	Hazwani Ismail, Wan Zul Haikal Hafiz Wan Zukiman	1
2.	STENTING IN POSTERIOR INFERIOR CEREBELLAR	ARTERY
	ANEURYSM: A CASE REPORT	
	Amelia Anggia, Ahmad Sobri Muda, Anas Tharek, Mohd Fandi Al Khafiz Ka	ımis, Rajeev
	Shamsuddin Perisamy, Siow Li Ting, Badrisyah Idris	

## CHALLENGING CATHETER MANAGEMENT IN A RARE CASE OF MYELOMENINGOCELE WITH SITUS INVERSUS TOTALIS: A CASE REPORT

Hazwani Ismail<sup>1</sup>, Wan Zul Haikal Hafiz Wan Zukiman<sup>1\*</sup>

<sup>1</sup>Department of Medicine, Faculty of Medicine and Health Sciences, University Putra

Malaysia, 43400 Serdang, Selangor, Malaysia

#### \*Corresponding author:

Wan Zul Haikal Hafiz Wan Zukiman, Department of Medicine, Faculty of Medicine and Health Sciences, University Putra Malaysia, 43400 Serdang, Selangor, Malaysia.

Email: <a href="mailto:zulhaikal@upm.edu.my">zulhaikal@upm.edu.my</a>

DOI: https://doi.org/10.32896/cvns.v6n1.1-7 Received: 11.11.2023 Revised: 29.03.2024 Accepted: 30.03.2024 Published: 31.03.2024

#### ABSTRACT

Individuals diagnosed with situs inversus totalis and renal failure have distinctive difficulties, such as an increased tendency to vascular stenosis and thrombosis during dialysis as a result of changed vascular patterns. A 19-year-old woman with myelomeningocele and situs inversus totalis is diagnosed with end-stage kidney disease (ESKD) as a result of bladder dysfunction. Due to the inadequacy of her vascular access, she commenced hemodialysis with the right permanent catheter. Regrettably, she developed a central line-associated bloodstream infection caused by Methicillin-Susceptible Staphylococcus aureus (MSSA), leading to septic shock. A CT scan and echocardiography revealed 9-mm tricuspid vegetation located around the cavoatrial junction. After a six-week course of intravenous cloxacillin, she was discharged with a left superior vena cava catheter. Without appropriate treatment, vascular issues might have serious repercussions for ESKD patients. Situs inversus presents extra challenges. Putting a catheter in a dextrocardia patient requires special skills. They also have unique vascular anatomy that necessitates different management than typical ESKD patients. Thus, a proper catheter size and type and imaging modalities like ultrasonography or venography are needed to simplify insertion and reduce complications. Renal failure treatment is perplexed by the presence of situs inversus totalis.

Keywords: Situs Inversus Totalis, Dextrocardia, ESKD, permanent catheter, vascular stenosis

#### **INTRODUCTION**

Situs inversus totalis typically does not result in any significant harm to an individual; however, it can occur alongside other congenital abnormalities, including those affecting the cardiovascular system (ventricular septal defect, atrial septal defect, tetralogy of Fallot, transposition of great arteries), the respiratory system (bronchiectasis, paranasal sinus deformity), and the digestive system (such as anal atresia, duodenal atresia) (Vaid et al., 2015). Patients with situs inversus totalis may develop renal failure as a result of many post issues, such infectious as glomerulonephritis, IgA nephropathy, and cystic kidney disease. The occurrence of spinal dysraphism in conjunction with situs inversus totalis is extremely unusual, and to this day, only a handful of examples have been documented (Vaid et al., 2015). The commonest spina bifida is myelomeningocele. It can be associated hydrocephalus, Arnold with Chiari malformation. VACTERL (vertebral defects, anal atresia, cardiac defects. tracheo-esophageal fistula, renal anomalies, and limb abnormalities), and **OEIS** (Omphalocele-Exstrophy-Imperforate Anus-Spinal Defects). This case report delves into the complex medical journey of a 19-year-old girl who faced a multitude of complications stemming from this rare combination.

#### **CASE REPORT**

A 19-year-old girl with myelomeningocele and situs inversus totalis caused complications include leg paralysis, bladder dysfunction, scoliosis, hydrocephalus, and cognitive impairment. Shunt placement for hydrocephalus and scoliosis correction were needed for the patient. Her neurogenic bladder caused vesicoureteral reflux and recurring urinary tract infections, resulting in ESKD. Due to her ESKD, the patient started hemodialysis, however her small size prevented fistula development. Dextrocardia rendered catheter insertion difficult and uncomfortable for the patient.

The patient had repeated catheter exchanges due to catheter malfunction after catheter implantation.

After three months on a permanent catheter, the patient came to the HDU for dialysis session with fever, cough, and shortness of breath intradialysis. The patient was not in respiratory distress, and the tunnelled catheter exit site discharged little pus. A CT scan revealed situs inversus, big impacted faeces in the rectum, focal fat stranding in the right paracolic gutter in the omentum, mild bilateral hydroureteronephrosis with diffusely enhanced urothelium, and several osseous lytic lesions, most likely brown tumours.

Based on the patient's clinical presentation, the patient was diagnosed with septic shock with a possible CLABSI (central line bloodstream infection). Urgent recommendations were made to arrange for a three-hour HD session through the available tunnelled catheter, remove it immediately after, and keep the patient in line-holiday for 24-48 hours. Additionally, arranging for Quinton line insertion through femoral vein for the next session.

Despite being on long-term antibiotics, the patient did not show signs of recovery, and echocardiogram was done, revealing a 9multilobulated hypodense mm lesion attached tricuspid to the valve posterior/septal leaflet, likely representing vegetation. Furthermore, an ill-defined hypodensity at the cavoatrial junction was causing mild luminal narrowing, likely representing thrombus formation. Bilateral subpleural consolidative densities with areas of central lucency were highly suggestive of septic emboli. Her blood methicillin-sensitive culture shows Staphylococcus aureus and requires a 6week antibiotic course. Repeated ECHO shows resolution of tricuspid vegetation. She was temporarily dialyzed via a femoral catheter during her hospital stay. She was discharged well at the opposite site (Left) of permanent catheter insertion and is awaiting a vascular appointment for fistula creation.

#### DISCUSSION

This case illustrates the complex history of a 19-year-old girl with several congenital abnormalities and multiple complications requiring ongoing medical management and surgical interventions. She has myelomeningocele, which is a subset of spina bifida. Spina bifida, which is the most common congenital abnormality of the central nervous system and involves a problem with neural tube closure. It may be classified anatomically into open spina bifida and closed spina bifida, with the former including myelomeningocele and myelocele as the most common forms.

A spinal cord segment protruding through a backbone gap causes myelomeningocele, which causes neurological impairments and other issues (Ntimbani et al., 2019). Myelomeningocele can induce limb muscle weakness or paralysis, bowel and bladder dysfunction, scoliosis, and cognitive deficits. requiring constant medical management and а multidisciplinary approach. Damage to the nerves that regulate the bladder muscles and sphincters can cause urine incontinence, UTIs, and other issues. Additionally, urinary tract vesicoureteral infections. reflux. and obstructive uropathy can cause end-stage kidney disease. Treatment aims to improve function and quality of life while reducing consequences and medical issues, including emotional and social challenges. Conventional postnatal surgical repair of a myelomeningocele defect is performed as soon as practically feasible after the exclusion of life-threatening associations, ideally during the first 48 hours of birth, to limit the danger of infection, chronic cerebrospinal fluid leakage, and for convenience of care.

Besides that, she has situs inversus, which is a rare genetic condition in which the organs in a person's chest and abdomen develop in a right-left formation instead of the normal left-right formation. Situs inversus can be associated with cystic kidney diseases and chronic renal failure. However, only three genes have been found to be responsible for some cases of cystic kidney diseases in situs inversus patients (Once et al., 2013). Patients with situs inversus who have chronic renal failure may have symptoms such as pallor and other signs of renal insufficiency (Mittal et al., 1994). Imaging tests such as an ultrasound, X-ray, CT scan, or MRI scan can be used to diagnose situs inversus and chronic kidney disease. If complications related to another defect or condition occur, these can be treated. It is important to inform healthcare providers if you have situs inversus so that future ailments can be diagnosed accurately due to the mirrored anatomy.

On the other hand, there are few types of dextrocardia that can be identified, which include dextrocardia with situs solitus, with situs inversus, and with situs ambiguous. Mirror-image Dextrocardia with situs inversus, inverted great arteries, and L-loop ventricles is the result of situs inversus with concordant L-bulboventricular loop, which is the opposite of normal cardiac anatomy (Maldjian et al., 2007). Dextrocardia occurs the migration of the due to Lbulboventricular loop's apex into the right hemithorax, which is the most common type of dextrocardia in the general population, occurring in one or two out of every 20,000 individuals .

In dextrocardia, the positions of the vessels that enter and exit the heart are reversed, with the aorta arising from the right ventricle and the pulmonary artery arising from the left ventricle, contrary to the usual positioning. This anomaly also extends to the superior vena cava, located left of the midline, and the inferior vena cava, situated to the right of the midline, in contrast to their normal anatomical positions (Maldjian et al., 2007). Furthermore, the coronary arteries responsible for supplying blood to the heart muscle are reversed in dextrocardia, with the right coronary artery arising from the left side of the aorta and the left coronary artery arising from the right side of the aorta, which differs from typical cardiac anatomy. These unique anatomical differences in dextrocardia hold significant implications for the diagnosis and medical treatment of affected individuals. Due to this condition, catheter insertion is more difficult because most catheters are designed for people with normal anatomy. Guided bv the KDOQI 2020 recommendations, catheter insertion should prioritize vascular access sites in the following order: right internal jugular, left internal jugular, external jugular, femoral, subclavian, and lumbar veins, to minimize procedural complications (Lok et al., 2019). However, in cases of situs inversus, where anatomy is mirrored, the preferred site for catheter placement should be the left internal jugular vein to accommodate the reversed anatomical landmarks.

Insertion of a catheter in a dextrocardia patient can be challenging, and multiple attempts may be necessary before successful placement. However, multiple attempts can also increase the risk of complications such as bleeding, infection, and damage to surrounding structures. To minimise the risk of complications during insertion. it's important catheter to implement catheter care bundle standard of procedure, select an appropriate catheter size and type, and ensure proper placement and securement of the catheter. Imaging plays a crucial role in identifying vessels for catheter insertion in patients with dextrocardia who require hemodialysis. Imaging techniques such as ultrasound or venography can help locate suitable vessels for catheter insertion and guide the placement of the catheter. Additionally, this can help to minimise the risk of complications and improve patient outcomes.

Furthermore, the synergistic collaboration amongst the nephrology, vascular, and interventional radiology teams is paramount. A unified approach and ongoing dialogue within this multidisciplinary team, aligned with the principles of the 'End-Stage Kidney Disease Life Plan,' serve not merely to preclude dialysis access-related complications but also to enhance overall patient morbidity and mortality outcomes. This holistic strategy underscores the necessity of a coordinated care paradigm, ensuring that each catheter insertion is not only technically successful but also aligned with the patient's broader treatment objectives and quality of life considerations.

## **REFERENCES:**

- Vaid, V., Kalra, S., Mahapatra, A., & Kumar, R. (2015). Split cord malformation and meningomyelocele with situs inversus totalis in a child: A rare association. Journal of Pediatric Neurology, 06(03), 269–272. doi:10.1055/s-0035-1557465
- Ntimbani, J., Kelly, A., & Lekgwara, P. (2019). Myelomeningocele - A literature review. Interdisciplinary Neurosurgery, 100502. doi:10.1016/j.inat.2019.100502
- Maldjian PD, Saric M. Approach to dextrocardia in adults: review. AJR Am J Roentgenol. 2007 Jun;188(6 Suppl):S39-49; quiz S35-8. doi: 10.2214/AJR.06.1179. PMID: 17515336.
- Mittal, R., Saxena, S., Hotchandani, R., Agarwal, S.K., Tiwari, S.C., & Dash, S.C. (1994). Situs inversus: a rare extrarenal association of Alport's syndrome. Nephron, 68 3, 390.
- 5. Onoe T, Konoshita T, Tsuneyama K, Hamano R, Mizushima I, Kakuchi Y, Yamada K, Hayashi K, Kuroda M, Kagitani S, Nomura H, Yamagishi M, Kawano M. Situs inversus and cystic kidney disease: Two adult patients with this Heterogeneous syndrome. Am J Case Rep. 2013;14:20-5. 10.12659/AJCR.883751. doi: 2013 28. Epub Jan PMID:

23569556; PMC3619039. PMCID:

 Lok CE, Huber TS, Lee T, Shenoy S, Yevzlin AS, Abreo K, Allon M, Asif A, Astor BC, Glickman MH, Graham J, Moist LM, Rajan DK, Roberts C, Vachharajani TJ, Valentini RP; National Kidney Foundation. KDOQI Clinical Practice Guideline for Vascular Access: 2019 Update. Am J Kidney Dis. 2020 Apr;75(4 Suppl 2):S1-S164. doi: 10.1053/j.ajkd.2019.12.001. Epub 2020 Mar 12. Erratum in: Am J Kidney Dis. 2021 Apr;77(4):551. PMID: 32778223.

## FIGURE LEGENDS:



Figure 1: CXR showing dextrocardia and a left permanent catheter in situ.



Figure 2: CT scan showing the location of the permanent catheter at IJV.

# STENTING IN POSTERIOR INFERIOR CEREBELLAR ARTERY ANEURYSM: A CASE REPORT

Amelia Anggia<sup>1</sup>, Ahmad Sobri Muda<sup>2\*</sup>, Anas Tharek<sup>2</sup>, Mohd Fandi Al Khafiz

Kamis<sup>2</sup>, Rajeev Shamsuddin Perisamy<sup>3</sup>, Siow Li Ting<sup>4</sup>, Badrisyah Idris<sup>5</sup>

<sup>1</sup>Radiology Departement, University of Brawijaya, Malang, Indonesia

<sup>2</sup>Radiology Department, Hospital Pengajar University Putra Malaysia, Selangor, Malaysia

<sup>3</sup>Department of Radiology, Kulliyyah of Medicine, International Islamic University Malaysia, Kuantan, Pahang, Malaysia

<sup>4</sup>Anaesthesiology and Critical Care Departement, Hospital Pengajar University Putra Malaysia, Selangor, Malaysia

<sup>5</sup>Neurosurgery Department, MSU Medical Centre, Selangor, Malaysia

#### \*Corresponding author:

Ahmad Sobri Muda, Radiology Department, Hospital Pengajar University Putra Malaysia, Selangor, Malaysia.

Email: <u>asobri@upm.edu.my</u>

DOI: https://doi.org/10.32896/cvns.v6n1.8-14 Received: 12.03.2024 Revised: 27.03.2024 Accepted: 30.03.2024 Published: 31.03.2024

#### ABSTRACT

Posterior inferior cerebellar artery (PICA) distal aneurysm is a relatively rare condition, accounting for 1% or less of all cerebral aneurysms. They may manifest as severe headaches, altered consciousness, lower cranial nerve dysfunction, brainstem compression, or posterior fossa symptoms such as nausea, vomiting, imbalance, or ataxia. Digital Subtraction Angiography (DSA) is the gold standard for detecting aneurysmal SAH. Endovascular treatment (EVT) has become the primary strategy for managing PICA aneurysms in recent years.

Keywords: Posterior Inferior Cerebellar Artery, Aneurysms, Stenting

#### **INTRODUCTION**

Aneurysms of the posterior inferior cerebellar artery (PICA) are uncommon. Most PICA aneurysms originate from the PICA-vertebral artery junction. Posterior circulation aneurysms, including PICA, has re-bleeding rates as high as 78%, primarily due to their relatively thin aneurysm wall and dissecting nature (Deora et a., 2020).

The most frequent clinical presentation of VA or PICA aneurysms is subarachnoid haemorrhage (SAH). Patients may experience headaches, and in a majority, vertigo and drowsiness. Additionally, unruptured VA, PICA, or vertebrobasilar aneurysms can lead to lower cranial nerve dysfunction, brainstem compression, or posterior fossa symptoms such as nausea, vomiting, imbalance, or ataxia (Hung, 2012).

The microsurgical approach is limited by anatomical corridors of the brainstem, petrous occipital bone, and multiple neurovascular structures occupying the cerebellomedullary and cerebellopontine cisterns. We present a case of ruptured PICA aneurysm treated with stenting.

#### CASE REPORT

A 59-year-old man with a Glasgow Coma Scale (GCS) of E1VTM1 (2T/15) and a blood pressure of 134/78 mmHg was transferred for endovascular treatment. Right external ventricular drainage done under general anaesthesia at a previous hospital. Upon arrival, the patient complained of neck stiffness, severe headache. blurred vision. vomiting. unsteady gait, and а decrease in consciousness from a GCS of 15 to 13. Plain brain CT scans revealed presence of SAH and intraventricular haemorrhages in the third and fourth ventricles, (Figure 1). angiography identified Cerebral an aneurysm of the right posterior inferior cerebellar artery (PICA) (Figure 2). The patient received an antiplatelet combination of aspirin 150 mg and clopidogrel 75 mg before the intervention. After MDT discussion, patient was decided for

endovascular treatment. Braided stent LEO+ Baby (Balt International) 2.0 mm x 12 mm navigated crossed the PICA aneurysm using microcatheter VASCOT (Balt International) 017 system (Figure 3.1). The stent fully developed with pattern flow of parent artery. There is constant stagnation following the stent placement (Figure 3B).

## DISCUSSION

Fusiform nonsaccular aneurysms at vertebrobasilar locations, more prevalent than in the anterior circulation due to dissections and carry a high risk of bleeding. Early treatment of ruptured aneurysms is crucial due to the high rebleeding rate and the potential risk of direct brainstem compression, leading to a high mortality rate (Hung 2012).

PICA aneurysms exhibit diversity in location and morphology, making it challenging to apply a single treatment modality consistently. The choice of treatment depends on factors such as aneurysm morphology, location, and the preference of the surgeon or interventionist. When considering aneurysm location, PICA aneurysms can be classified as proximal (including those arising at the origin of the PICA-vertebral junction or within the anteromedullary segment) and distal (located after the anteromedullary segment) (Khayat et al., 2020).

Most PICA aneurysms arise at the junction of the PICA and vertebral artery, with approximately one-third arising more distally. The variable origin, tortuous course, and close proximity to vital neurovascular structures present challenges for both endovascular and open surgical approaches (Lheto et al., 2014).

Endovascular treatment (EVT) has become the predominant strategy for treating PICA aneurysms recently, accounting for 55.4% of ruptured PICA aneurysms. EVT reduces the risk of direct brainstem injury and anaesthesia-related complications, making it preferable for patients with poor clinical status. EVT strategies such as stand-alone coiling, stent-assisted coiling, direct coiling, flow diversion, and parent artery occlusion with coils have been successfully used to treat PICA aneurysms. EVT-related complications, including intraprocedural ruptures and new-onset infarcts, are not rare, with reported morbidity from EVTrelated imaging infarcts reaching up to 33%, although many of them may not show apparent symptoms (Chen et al., 2018).

Patients who survived the hospital admission period were scheduled for clinical follow-up in the outpatient clinic at 6 weeks and for angiographic follow-up at 6 and 18 months. In some patients treated with coil occlusion of the aneurysm including the PICA origin and/or vertebral artery occlusion, only MR imaging and MR angiography (MRA) follow-up was performed (Peluso and Rooji, 2008).

#### CONCLUSION

PICA aneurysms present challenging lesions associated with high mortality rates if early diagnosis is missed. The primary goals of examination are to identify the site of the vascular lesion and determine whether immediate intervention is required. CT and MRI serve as the imaging modalities of choice for identifying and locating the infarct and detecting vascular area abnormalities. Digital Subtraction Angiography (DSA) stands as the gold standard for detecting aneurysmal SAH (Andrew et al., 2023).

Endovascular treatment (EVT) has emerged as the dominant strategy for managing PICA aneurysms, as it reduces the risk of direct brainstem injury and anaesthesia-related complications, particularly in patients with poor clinical status.

Involvement of a multidisciplinary team is crucial for diligent treatment planning, given the complexity of these lesions. The information provided in this report is intended to assist in the decision-making process.

## DATA AVAILABILITY STATEMENT

https://interventionjournal.padimedical.c om/external/view/421a04a0-9172c6b6-64725c36-ac89bfd6-2d85d9d0/

## **REFERENCES:**

Harsh Deora, Nitish Nayak,
Priyadarshi Dixit et all. Surgical
Management and Outcomes of
Aneurysms of Posterior Inferior
Cerebellar Artery: Location-Based
Approaches with Review of Literature. J
Neurosci Rural Pract. 2020 Jan; 11(1):
34–43.

2. Khayat HA, Hawkes CM and Algird AR. Case Report: Endovascular Treatment of a Giant Distal PICA Aneurysm in Association with a Cerebellar AVM: A Report on Treatment Considerations and a Literature Review. Front. Neurol. 11:611377. doi: 10.3389/fneur.2020.611377

3. Lheto H, Harati A, Niemelä M, Kurosaki Y, Chin M, Yamagata S, et al. Distal posterior inferior cerebellar artery aneurysms: clinical features and outcome of 80 patients. World Neurosurg. (2014) 82:702–13. doi: 10.1016/j.wneu.2014.06.012

4. Chen X, Sun Z, Shi L, Xu L, Yu J, Fang B. et al. Endovascular management of ruptured distal posterior inferior cerebellar artery aneurysms: A retrospective cohort study. Medicine (Baltimore) 2018;97(49 e):13300.

5. J.P Peluso, W.J Van Rooji. Posterior Inferior Cerebellar Artery Aneurysms: Incidence, Clinical Presentation, and Outcome of Endovascular Treatment. American Journal of Neuroradiology January 2008, 29 (1) 86-90

6. Andrew, M. et all. Cerebral Aneurysm. StatPearls Publishing;2023. Available from: Cerebral Aneurysm -StatPearls - NCBI Bookshelf (nih.gov)  Che Wei Hung. Posterior Inferior Cerebellar Artery Aneurysm mimicking cervical myeloradiculopathy. 2012.
Formosan Journal of Surgery. Volume 45, Issue 3, June 2012, Pages 88-92

## FIGURE LEGENDS:



Figure 1: CT scan showed intraventricular haemorrhages at third and fourth ventricle causing obstructive hydrocephalus and subarachnoid haemorrhage.



Figure 2: Cerebral angiogram showed aneurysm of the right PICA.



Figure 3: Stenting placement (A), stenting procedure (B), before (C) and post stenting (D).